

Not Every Finite Lattice is Embeddable in the Recursively Enumerable Degrees*

A. H. LACHLAN[†] AND R. I. SOARE[‡]

[†] *Department of Mathematics, Simon Fraser University, Burnaby 2,
British Columbia, Canada; and*

[‡] *Department of Mathematics, University of Chicago, Chicago, Illinois 60637*

A certain lattice with eight elements is shown to be not embeddable as a lattice in the recursively enumerable degrees. This refutes the well-known Embedding Conjecture which asserted that every finite lattice could be so embedded.

Since the papers of Kleene, Post, and Spector [2, 10, 20], a great deal of important research has been devoted to the degrees \mathcal{D} of unsolvability, particularly their elementary theory, $\text{Th}(\mathcal{D})$ and their algebraic structure as an upper semi-lattice. Lachlan [4] proved undecidability of $\text{Th}(\mathcal{D})$ by proving that all finite lattices could be embedded as initial segments of degrees. Simpson [18] later obtained a more exact classification by showing $\text{Th}(\mathcal{D})$ to be recursively isomorphic to the truth set of second order arithmetic. Building on a technique of Lerman [7], Lachlan and Lebeuf [6] showed that every countable upper semi-lattice could be embedded in \mathcal{D} as an initial segment, and Nerode and Shore [9] used this to obtain a simple proof of Simpson's result.

Among classes of degrees, those containing recursively enumerable sets (the r.e. degrees) have received particular attention since the time of Post [10] partly because of the widespread occurrence of r.e. sets in algebra and number theory as explained in [19], and their application in famous undecidability results. Both the decision problem for the elementary theory and embedding questions are much more difficult for the r.e. degrees \mathcal{R} than for \mathcal{D} . For example, Friedberg [1] and Muchnik [8] invented the priority method just to prove that there are more than two r.e. degrees, and the proof that \mathcal{R} is not a lattice was only given in [3]. Sacks [14] showed that the r.e. degrees are dense, probably the most pleasing property of \mathcal{R} to have been proved. Many other elementary properties of \mathcal{R} have been proved, some by the exercise of great ingenuity, but the overall picture has remained obscure. A survey of the literature can be found in [19].

Here we are chiefly concerned with what finite lattices can be embedded in \mathcal{R} as lattices, i.e., we are insisting that both joins and meets be preserved by


* This research was partially supported by NSF Grant MCS 76-07033 A01.

the embedding. The first result along this line, due independently to Lachlan [3] and Yates [22], showed that the lattice



can be embedded in \mathcal{R} as a lattice with least element mapped to 0. Building on this Thomason [21] and Lachlan [5], and independently, Lerman (unpublished) showed that all countable distributive lattices can be embedded in \mathcal{R} as lattices. Using a much more complex technique of enumeration Lachlan [5] showed how the two 5-element nondistributive lattices can be embedded in \mathcal{R} as lattices. This led many to attempt to prove the Embedding Conjecture [11, 16, 19] which asserts that every finite lattice can be embedded in \mathcal{R} as a lattice. A solution of this conjecture is necessary to decide even which *existential* sentences are true in \mathcal{R} .

In this paper we establish that the lattice \mathcal{L} (also called S_8) with diagram



is not embeddable in \mathcal{R} as a lattice. Lerman was the first to suggest that the Embedding Conjecture might be false and in particular that S_8 might not be embeddable. Although great difficulties remain, we hope this will at least point the way to a solution of the whole embedding problem for \mathcal{R} , and will thus shed some light on the decision problem for its elementary theory.

Preliminaries. Below we shall deal with sets, functions, and functionals which are enumerated in ω stages. We think of all these enumerations as occurring simultaneously. For brevity we deliberately employ an ambiguous notation. If A denotes an r.e. set then it also denotes the particular enumeration of the r.e. set, and within the context of a stage denotes the current approximation to A . If the stage is not clear then the approximation to A at stage s , i.e., the finite set enumerated in A before stage s , is denoted $A[s]$. The sequence $\langle A[s]; s < \omega \rangle$ is increasing and strongly r.e. Finally, A also denotes the characteristic function of A . If Ψ denotes a p.r. functional Ψ also denotes the enumeration of Ψ : $\langle \Psi[s]; s < \omega \rangle$ a strongly r.e. increasing sequence of finite functionals. In the context of stage s , Ψ refers to $\Psi[s]$ the current approximation.

For functions our terminology is unorthodox. Functions have natural-number arguments but are not necessarily total. If φ is a function φ also denotes

the particular enumeration of it with which we are concerned. But in this case $\langle \varphi[s]: s < \omega \rangle$ is an r.e. sequence of p.r. functions such that $x \in \text{dom } \varphi[y]$ is a recursive binary relation and for all x, y

$$\varphi(x) = y \leftrightarrow \exists t \forall s (s \geq t \rightarrow \varphi(x)[s] = y).$$

We are not requiring that $\varphi[s]$ be finite nor that $\varphi[s+1]$ extend $\varphi[s]$. Clearly, the total functions enumerable in this sense are those of degree $\leq 0'$. If A is an r.e. set and φ is a function, then φ *respects* A means that for all x, s

$$\begin{aligned} &\varphi(x)[s] \text{ defined \& } (\varphi(x)[s+1] \text{ undefined or } \varphi(x)[s+1] \neq \varphi(x)[s]) \\ &\rightarrow \exists y (y < \varphi(x)[s] \& y \in A[s+1] - A[s]). \end{aligned}$$

For any functional Ψ and set A we define the *use function* ψ of Ψ with respect to A by:

$$\begin{aligned} \psi(x)[s] = \max(\{ \mu y (\Psi(A \upharpoonright y)(x)[s] \text{ is defined}) \cup \{x\} \\ \cup \{ \psi(x)[t]: t < s \& \psi(x)[t] \text{ is defined} \}). \end{aligned}$$

Note that ψ respects A and that ψ is total if $\Psi(A)$ is total.

For any set A its degree will be denoted \mathbf{a} , i.e., by the corresponding lower case letter.

We now move directly to the proof of

THEOREM. \mathcal{L} cannot be embedded as a lattice in the upper semilattice of r.e. degrees.

It suffices to prove something apparently weaker, namely, that


$$\mathcal{L} \text{ cannot be embedded with its least element } 0_{\mathcal{L}} \text{ being mapped to } 0. \quad (1)$$

From the latter by relativization for any r.e. degree \mathbf{b} there is no embedding of \mathcal{L} as a lattice into the \mathbf{b} -r.e. degrees with $0_{\mathcal{L}}$ being mapped to \mathbf{b} . From [2, p. 568] we have:

LEMMA 1. *Let $\mathbf{a}_0, \mathbf{a}_1$ be r.e. degrees and \mathbf{b} a degree not necessarily r.e. such that $\mathbf{b} \leq \mathbf{a}_0, \mathbf{a}_1$; then there exists an r.e. degree $\mathbf{c} \leq \mathbf{a}_0, \mathbf{a}_1$ such that $\mathbf{b} \leq \mathbf{c}$.*

An immediate consequence is that an embedding of \mathcal{L} as a lattice into the r.e. degrees with $0_{\mathcal{L}}$ going to \mathbf{b} would simultaneously be an embedding into the \mathbf{b} -r.e. degrees. Thus below we need only prove (1).

For proof by contradiction, suppose given r.e. sets A^0, A^1, C^0, C^1, C^2 which witness the embedding of the lattice \mathcal{L} in the upper semilattice of r.e. degrees in the manner of the diagram:



It is convenient to suppose that C^i contains only even numbers congruent to $i \pmod{3}$. Let $C = \bigcup \{C^i: i < 3\}$ then C contains only even numbers and we may assume that A_0 and A_1 agree with C on the even numbers. For $\{i, j, k\} = \{0, 1, 2\}$ let τ^i be the enumeration of a function witnessing $c^i \leq c^j \cup c^k$ in the sense that

- (i) $\tau^i[s]$ is total and converges as $s \rightarrow \infty$;
- (ii) τ^i respects $C^j \cup C^k$;
- (iii) for all x and s , $\tau^i(x)[s]$ is increasing in x and nondecreasing in s , and $x < \tau^i(x)[s]$;
- (iv) for all x, s

$$x \in C^i[s+1]$$

$$- C^i[s] \rightarrow \exists y (y < \tau^i(x)[s] \text{ \& } y \in C^j[s+1] - C^j[s] \vee y \in C^k[s+1] - C^k[s]).$$

Let $\tau(x) = \max\{\tau^i(x): i < 3\}$, then τ respects C .

To verify the existence of the enumerations $\tau^i[s]$ and $C^i[s]$, fix recursive functionals Φ^i such that $C^i = \Phi^i(C^j \cup C^k)$, and let φ_i be the use function for Φ_i . Let $\hat{C}^i[s]$, $s \in \omega$, be any recursive enumeration of C^i , and define recursive functions

$$l^i(s) = (\mu x)(\forall y < x)[\hat{C}^i[s] = \Phi^i(\hat{C}^j \cup \hat{C}^k)[s]].$$

Choose stages $t_0 < t_1 < \dots$ such that

$$(\forall i < 3)[\hat{C}^i[t_{n+1}] - \hat{C}^i[t_n] \neq \emptyset \quad \text{and} \quad n \leq l^i(t_n)].$$

Let $z_n^i = \mu x [x \in \hat{C}^i[t_{n+1}] - \hat{C}^i[t_n]]$. Define $C^i[s] = \hat{C}^i[t_s]$, $g(i) = (i+1) \pmod{3}$, and

$$\begin{aligned} \sigma^i(x)[s] &= \varphi_i(x)[t_s] & \text{if } x \leq s \\ &= 1 + z_s^{g(i)} & \text{if } x > s. \end{aligned}$$

Define $\tau^i(x)[s] = \max\{1 + x, \sigma^i(x)[s], \tau^i(x)[s - 1]\}$. Clearly, these enumerations satisfy (i), (ii), and (iii) above. For (iv), assume $x \in C^i[s + 1] - C^i[s]$. If $x > s$, then

$$\tau^i(x)[s] > z_s^{g(i)} \in (C^j \cup C^k)[s + 1] - (C^j \cup C^k)[s].$$

If $x < s$, then $\tau^i(x)[s] \geq \varphi^i(x)[s]$. But $x \in \hat{C}^i[t_{s+1}] - \hat{C}^i[t_s]$, $l^i(t_s) \geq s \geq x$, and $l^i(t_{s+1}) \geq s + 1 \geq x$ imply there is some $y < \varphi^i(x)[s]$, such that

$$y \in (\hat{C}^j \cup \hat{C}^k)[t_{s+1}] - (\hat{C}^j \cup \hat{C}^k)[t_s].$$

Let θ^0, θ^1, ψ be enumerations of functions. With respect to this triple of enumerations we say n is *active at stage* s if $\theta^0(n)$, $\theta^1(n)$, and $\psi(n)$ are all defined and

$$\tau\psi(n) < \theta^0(n) = \theta^1(n).$$

Call n *permanently active* if, for some s , n is active all stages $\geq s$.

LEMMA 2. *Given a recursive enumeration ψ of a use function which respects C we can effectively enumerate, uniformly in C , total functions θ^0, θ^1 , respecting A^0, A^1 , respectively, such that if ψ is total then there are infinitely many n which are permanently active. Further, for all n and s*

$$(n \text{ active at stage } s \text{ but not at stage } s + 1) \rightarrow \\ \exists x(x < \psi(n)[s] \ \& \ x \in C[s + 1] - C[s]).$$

Proof. Define $\theta^0(x)[0] = \theta^1(x)[0] = x$. If $i < 2$, $s > 0$, $\psi(x)[s - 1] = \psi(x)[s]$ is defined, $\theta^i(x)[s - 1] > \tau\psi(x)[s - 1]$, and $\tau\psi(x)[s] > \tau\psi(x)[s - 1]$, then let

$$\theta^i(x)[s] = \tau\psi(x)[s] + 1.$$

If $s > 0$, $\psi(x)[s - 1] = \psi(x)[s]$ is defined, $\theta^0(x)[s - 1] \leq \tau\psi(x)[s]$, and $(A^0[s] - A^0[s - 1]) \cap x \neq \emptyset$, let

$$\theta^0(x)[s] = \tau\psi(x)[s] + 1.$$

If $s > 0$, $\psi(x)[s - 1] = \psi(x)[s]$ is defined, $\theta^0(x)[s - 1] > \tau\psi(x)[s]$, and $(A^1[s] - A^1[s - 1]) \cap x \neq \emptyset$, let

$$\theta^1(x)[s] = \theta^0(x)[s - 1].$$

In all other cases if $i < 2$ and $s > 0$ let $\theta^i(x)[s] = \theta^i(x)[s - 1]$.

If $\tau\psi(x)[s] > \tau\psi(x)[s - 1]$ while $\psi(x)[s] = \psi(x)[s - 1]$ then $(C[s] - C[s - 1]) \cap \tau\psi(x)[s - 1] \neq \emptyset$ since τ respects C . From this it is clear that θ^0, θ^1 respect A^0, A^1 , respectively.

Observe that if at some stage $\theta^i(x) > \tau\psi(x)$ then in fact $\theta^i(x) = \tau\psi(x) + 1$ because $\psi(x)[s] \geq x$ and $\tau(y)[s] > y$, so $\theta^i(x) < \tau(\psi(x))$ at $s = 0$. Further, if $\theta^i(x)[s] = \tau\psi(x) + 1$ and $\theta^i(x)[t] = \tau\psi(x)[t] + 1$ with $s < t$, we cannot have $\psi(x)[t] > \psi(x)[s]$ unless some number $< x$ enters A^i at a stage v , $s \leq v \leq t$. For consider the least $u > s$ with $\psi(x)[u] > \psi(x)$ and let v be greatest such that $s \leq v < u$ and $\psi(x)[v]$ is defined, then

$$\tau\psi(x)[u] \geq \tau\psi(x)[v] + 1 \geq \theta^i(x)[v] = \theta^i(x)[u].$$

Since $\tau\psi(x)[w]$ is nondecreasing in w from the definition of θ^i , $\theta^i(x)[t] > \tau\psi(x)[t]$ implies that some number $< x$ enters A^i at a stage $\geq u$ and $< t$. The above observation together with the convergence of τ guarantees the convergence of θ^i for $i < 2$.

Suppose ψ is total. For infinitely many n there exists s such that at stage s , $\psi(n)$ is defined, C has converged on $\psi(n)$, but A^0 has not yet converged on n . Otherwise, A^0 would be recursive in C . Thus using an A^0 -oracle we can effectively enumerate an infinite sequence $\langle (n(i), s(i)) : i < \omega \rangle$ such that for all i , $n(i) < n(i+1)$, $C[s(i)]$ and C agree on $\psi(n(i))$, $A^0[s(i)]$ and A^0 agree on $n(i)$, and $\theta^0(n(i))[s(i)] > \tau\psi(n(i))[s(i)]$. Clearly, $\psi(n(i))[s] = \psi(n(i))[s(i)]$ for all $s \geq s(i)$ because ψ respects C , whence also $\theta^0(n(i))[s] > \tau\psi(n(i))[s]$ for all $s \geq s(i)$. There are infinitely many i such that at stage $s(i)$, A^1 has not yet converged on $n(i)$. Otherwise, A^1 would be recursive in A^0 . If $x < n(i)$ is enumerated in A^1 at stage $t-1 \geq s(i)$ then by definition of θ^1 , $\theta^1(n(i))[t] = \theta^0(n(i))[t]$. Further, $\theta^1(n(i))[s] = \theta^0(n(i))[t]$ for all $s \geq t$. Thus $n(i)$ is permanently active. This completes the proof of the lemma.

Let Ψ be a recursive enumeration of a functional and ψ be the use function of Ψ with respect to C . Then ψ respects C . Let θ^0, θ^1 be the enumerations obtained by Lemma 2 from ψ . We effectively enumerate a function φ as follows:

$$\varphi(x)[0] = 0.$$

If n is active at stage s , $\Psi(C)(n)[s]$ is defined, and

$$\exists x(\psi(n) \leq x < \tau\psi(n) \ \& \ x \in C[s+1] - C[s]),$$

then $\varphi(n)[s+1] = 1 + \Psi(c)[n](s)$. Otherwise, $\varphi(n)[s+1] = \varphi(n)[s]$.

LEMMA 3. φ is total and is recursive in both A^0 and A^1 uniformly in Ψ, ψ .

Proof. Note from Lemma 2 that θ^0, θ^1 are recursive in A^0, A^1 , respectively, uniformly in ψ . For $i = 0, 1$, once $\theta^i(n)$ has settled down and all members $< \theta^i(n)$ have entered C , $\varphi(n)$ cannot change. This is enough.

Fix a simultaneous recursive enumeration $\langle (W_e^0, W_e^1, W_e^2) : e < \omega \rangle$ of all

triples of r.e. sets and a recursive partition of ω into infinite sets R_e^i , $(i, e) \in 3 \times \omega$. Members of R_e^i are called (i, e) -numbers.

With respect to the triple of enumerations θ^0, θ^1, ψ we define enumerations of r.e. sets G^0, G^1, G^2 as follows. Let $\{i, j, k\} = \{0, 1, 2\}$, then enumerate x in G^i at stage t if there exist e, s, t, n such that the following conditions hold:

- (1) x is an (i, e) -number, $x \in W_e^i[s]$;
- (2) $s < t$, $\tau\psi(n)[s] \leq x$;
- (3) n is active at all stages $\geq s$ and $< t$;
- (4) t is the least number $> s$ such that $(C[t] - C[s]) \cap \psi(n) \neq \emptyset$,

$$(C^j[t] - C^j[s]) \cap \tau\psi(n)[s] \neq \emptyset$$

and

$$(C^k[t] - C^k[s]) \cap \tau\psi(n)[s] \neq \emptyset.$$

LEMMA 4. *Let Ψ be an enumeration of a functional, ψ be the use function of Ψ with respect to C . Let θ^0, θ^1 be generated from ψ by Lemma 2 and then let φ and G^0, G^1, G^2 be defined as just described. One of the following three possibilities holds:*

- (P1) $\Psi(C)$ is not total.
- (P2) $\Psi(C)$ is total and $\Psi(C) \neq \varphi$.
- (P3) for all i, j, k such that $\{i, j, k\} = \{1, 2, 3\}$ G^i is recursive in both C^j and C^k ; further, one of G^0, G^1, G^2 is nonrecursive.

Proof. For argument by contradiction, suppose (P1), (P2), (P3) all fail. Let $\{i, j, k\} = \{1, 2, 3\}$. Suppose G^0, G^1, G^2 are all recursive. Then there exists e independent of i such that $W_e^i = \omega - G^i$ and no (i, e) -number is ever enumerated in G^i . Define $\lambda^i(s)$ to be the greatest (i, e) -number in $W_e^i[s]$ and let $\lambda(s) = \min\{\lambda^i(s) : i < 3\}$. Then $\lambda(s)$ is defined for all sufficiently large s is non-decreasing and unbounded as s increases. Note that $\lambda(s)$ is a recursive function.

Fix n and s such that $\tau\psi(n)[s] \leq \lambda(s)$ and n is active at stage s . Then n is active at all stages $\geq s$. Otherwise, let t be the first stage $> s$ at which n is not active. Then from Lemma 2

$$\exists x(x < \psi(n)[s] \ \& \ x \in C[t] - C[s]).$$

From this it follows that two of $C^0[t] - C^0[s]$, $C^1[t] - C^1[s]$, and $C^2[t] - C^2[s]$ intersect $\tau\psi(n)[s]$. Hence, for some $i < 3$, $\lambda^i(s)$ is enumerated in G^i making $G^i \cap W_e^i \neq \emptyset$, contradiction. It is also clear that if $\tau\psi(n)[s] \leq \lambda(s)$ and n is active at stage s then no number $< \psi(n)[s]$ is enumerated in C after stage s . Since $\psi(n) \geq n$ whenever $\psi(n)$ is defined and there are infinitely many permanently active n from Lemma 2, we have C recursive. Thus one of G^0, G^1, G^2 is nonrecursive.

Now we turn to the proof that G^i is recursive in both C^j and C^k . To this end, observe that there is a 4-ary recursive function π with the following property. If $u > s$, n is active at all stages $\geq s$ and $\leq u$, $y < \tau\psi(n)[s]$, $y \in C[u] - C[s]$, and

$$\neg \exists x(x < \psi(n)[s] \ \& \ x \in C[u] - C[s])$$

then $\pi(s, u, n, y)$ is the least $t > s$ such that

$$\exists x(x < \psi(n)[s] \ \& \ x \in C[t] - C[s]).$$

Such t must exist because otherwise $\varphi(n) \neq \Psi(C)(n)$ which would mean that one of (P1), (P2) holds. For other quadruples s, u, n, y let $\pi(s, u, n, y) = 0$. We can now show that G^i is recursive in C^j as follows. Given x find v such that $C^j[v]$ and C^j agree on x . Let

$$v' = \max\{\pi(s, v, n, y): y \in C^j[v] - C^j[s], s < \omega, n < \omega, y < \omega\}.$$

Since $\pi(s, v, n, y) = 0$ for $s \geq v$, and $\langle C^j[s]: s < \omega \rangle$ and $\langle \text{dom } \psi[s]: s < \omega \rangle$ are strongly r.e. sequences of finite sets, v' exists and may be effectively computed from v . Then $x \in G^i$ if and only if $x \in G^i[v']$ by clause (3) in the definition of G^i . This completes the proof that G^i is recursive in C^j and also the proof of Lemma 4.

Since (P3) is impossible from our assumption that $c^0 \cap c^1 = c^1 \cap c^2 = c^2 \cap c^0 = 0$, the argument given above shows that for any p.r. functional Ψ we can effectively enumerate uniformly in Ψ a function φ which is recursive in both A^0 and A^1 uniformly in Ψ . Further, $\varphi \neq \Psi(C)$. Let $\langle \Psi_i: i < \omega \rangle$ be a standard enumeration of the p.r. functionals and $x \mapsto ((x)_0, (x)_1)$ be a recursive bijection from ω to $\omega \times \omega$. Define $\delta(x) = \varphi_{(x)_0}(x)_1$, where φ_i comes from Ψ_i by our uniform method of enumerating φ from Ψ . Now δ is recursive in both A_0 and A_1 . Were δ recursive in C , there would be an r.e. sequence of p.r. functionals $\langle \Phi_i: i < \omega \rangle$ such that $\varphi_i = \Phi_i(C)$ for all $i < \omega$. But from the recursion theorem $\Phi_i = \Psi_i$ for some i which contradicts $\varphi_i \neq \Psi_i(C)$. Thus δ is not recursive in C and we have a degree $\mathbf{d} \leq \mathbf{a}^0, \mathbf{a}^1$ with $\mathbf{d} \not\leq \mathbf{c}$. From Lemma 1 $\mathbf{c} \neq \mathbf{a}^0 \cap \mathbf{a}^1$ in the upper semilattice of r.e. degrees which is the final contradiction we have been seeking.

REFERENCES

1. R. M. FRIEDBERG, Two recursively enumerable sets of incomparable degrees of unsolvability, *Proc. Nat. Acad. Sci. U.S.A.* **43** (1957), 236-238.
2. S. C. KLEENE AND E. L. POST, The upper semi-lattice of degrees of recursive unsolvability, *Ann. of Math. (2)* **59** (1954), 379-407.
3. A. H. LACHLAN, Lower bounds for pairs of r.e. degrees, *Proc. London Math. Soc.* (3) **16** (1966), 537-569.

4. A. H. LACHLAN, Distributive initial segments of the degrees of unsolvability, *Z. Math. Logik Grundlagen Math.* **14** (1968), 457-472.
5. A. H. LACHLAN, Embedding nondistributive lattices in the recursively enumerable degrees, in "Conference in Mathematical Logic, London 1970," pp. 149-177, Lecture Notes in Mathematics No. 255, Springer-Verlag, New York, 1972.
6. A. H. LACHLAN AND R. LEBEUF, Countable initial segments of the degrees of unsolvability, *J. Symbolic Logic* **41** (1976), 289-300.
7. M. LERMAN, Initial segments of the degrees of unsolvability, *Ann. of Math.* **93** (1971), 365-389.
8. A. A. MUCHNIK, On the unsolvability of the problem of reducibility in the theory of algorithms (Russian), *Dokl. Akad. Nauk SSSR* **108** (1956), 194-197.
9. A. NERODE AND R. A. SHORE, Second order logic and first order theories of reducibility orderings, to appear.
10. E. L. POST, Recursively enumerable sets of positive integers and their decision problems, *Bull. Amer. Math. Soc.* **50** (1944), 284-316.
11. R. W. ROBINSON, Interpolation and embedding in the recursively enumerable degrees, *Ann. of Math.* **93** (1971), 285-314.
13. G. E. SACKS, On the degrees less than $0'$, *Ann. of Math.* **77** (1963), 211-231.
14. G. E. SACKS, The recursively enumerable degrees are dense, *Ann. of Math.* (2) **80** (1964), 300-312.
15. G. E. SACKS, "Degrees of unsolvability," rev. ed., Annals of Mathematics Studies, No. 55, Princeton Univ. Press, Princeton, N.J., 1966.
16. J. R. SHOENFIELD, The decision problem for recursively enumerable degrees, *Bull. Amer. Math. Soc.* **81** (1975), 973-977.
17. S. G. SIMPSON, Degrees of unsolvability: A survey of results, in "Handbook of Mathematical Logic" (J. Barwise, Ed.) North-Holland, Amsterdam, 1977.
18. S. G. SIMPSON, First-order theory of the degrees of recursive unsolvability, *Ann. of Math.* (2) **105** (1977), 121-139.
19. R. I. SOARE, Recursively enumerable sets and degrees, *Bull. Amer. Math. Soc.* **84** (1978), 1149-1181.
20. C. SPECTOR, On degrees of recursive unsolvability, *Ann. of Math.* (2) **64** (1956), 581-592.
21. S. K. THOMASON, Sublattices of the recursively enumerable degrees, *Z. Math. Logik Grundlagen Math.* **17** (1971), 273-280.
22. C. E. M. YATES, A minimal pair of recursively enumerable degrees, *J. Symbolic Logic* **31** (1966), 159-168.